Vaccine Schedule for Your Dog

We make the following recommendations for maximal protection against certain serious infectious diseases that could affect your pet. New vaccines are now available and the age of the dog makes each series an individual one.

Distemper......A viral disease of the lungs, intestines and heartHepatitis......A viral disease of the liverLeptospirosisA bacterial disease of the urinary systemCorona......A viral disease of the intestinesParainfluenza......Infectious bronchitis (kennel cough)Parvo enteritis......A serious viral disease of the intestines and heartMeasles..........Used to protect young puppies against distemperRabies........A viral disease, fatal to man and animalsBordetella........Infections bronchitis (kennel cough – severe)Lyme Disease..........A bacterial disease transmitted by ticks

Program	Age	Vaccinate Now		Remainder of Series & Age Given	
1	6 wks.	DMP/PARVO	DHLPPC-10 wks.	DHLPPC/BORD-14 wks.	*DHLPPC/RABIES-18 wks.
2	7 wks.	DMP/PARVO	DHLPPC-10 wks.	DHLPPC/BORD-14 wks.	*DHLPPC/RABIES-18 wks.
3	8 wks.	DMP/PARVO	DHLPPC-11 wks.	DHLPPC/BORD-14 wks.	*DHLPPC/RABIES-18 wks.
4	9 wks.	DMP/PARVO	DHLPPC-12 wks.	DHLPPC/BORD-15 wks.	*DHLPPC/RABIES-18 wks.
5	10 wks.	DHLPPC	DHLPPC/BORD – 14wks.		*DHLPPC/RABIES-18 wks.
6	11 wks.	DHLPPC	DHLPPC/BORD – 14wks.		*DHLPPC/RABIES-18 wks.
7	12 wks.	DHLPPC	DHLPPC/BORD – 14wks.		*DHLPPC/RABIES-18 wks.
8	13 wks.	DHLPPC	DHLPPC/BORD – 14wks.		*DHLPPC/RABIES-19 wks.
9	14 wks.	DHLPPC/BORD	DHLPPC/RABIES – 18 wks.		
10	15 wks.	DHLPPC/BORD	DHLPPC/RABIES – 18 wks.		6 Month
11	16 wks.	DHLPPC/BORD	DHLPPC/RABIES – 19 wks.		Neuter
12	OLDER	DHLPPC/BORD	DHLPPC/RABIES – in 3 wks.		Microchip

PUPPIES

*Rottweiller and Doberman breeds should be blood tested at 4 weeks after last vaccine to see if adequate antibody levels for Parvo exist.

DHLPPC.....Yearly Booster Bordetella....Yearly Booster Rabies (under 1 year of age).....Yearly Booster Rabies (over 1 year of age).....Three-Year Booster Lyme (for dogs at risk).....Yearly Booster

ALL HOSPITALIZED DOGS MUST HAVE CURRENT DHLPPC AND RABIES VACCINATIONS

Bordetella intranasal vaccine lasts 12 months and is recommended for all dogs. It helps significantly reduce the incidence of severe upper respiratory infections (kennel cough).

Parasite Alert

Internal parasite tests are recommended for all puppies, pets with diarrhea, and at yearly examinations. Please bring a fresh specimen of your pet's stool (feces) for this important test. Remember, most parasites are not visible to the naked eye and only a microscopic exam will reveal the presence of the eggs or larvae.

PREPAID FECAL PROGRAM

Give worming medicine by mouth today and repeat every 2 weeks until gone. Drop off fresh stool sample 3 weeks after the last dose.